


INVERTEBRATES

FINAL PROJECT

BY GEMMA SHEPHERD



ECHINODERMATA

Sea cucumber

RIGIDITY & FLEXIBILITY
Sea cucumbers have mutable collagenous tissues which can extensively change their mechanical properties. Their skin can switch between soft, normal & rigid states in the blink of an eye. Their skin contains collagen fibrils which are able to bind strongly or not with other substances around them, switching from an interlinked rigid state to a loose flexible state.

SMART POLYMER
Researchers studying sea cucumber skin have developed a polymer which is flexible in aqueous solvent & becomes rigid when the solvent evaporates. They hope to use this for 'self-healing' medical devices & responsive brain implants.

SOFT ← NORMAL → HARD
- Hydrogen bonding inhibited
- H bonding activated

Electrical impulses

CNIDARIAN

Stony corals

CORAL GROWTH
Polyps secrete skeletons of calcium carbonate (CaCO_3) in the form of a cup in which the polyps sit. Occasionally, the polyps lift up & create a new base above it, elevating the coral upward using carbon dioxide & calcium from the ocean.

CARBON SEQUESTERING CEMENT
Mimicking the mechanism by which corals build their skeletons could transform the cement industry, which is currently responsible for 8% of total CO_2 emissions. The cement company Calceira now makes cement by dissolving waste CO_2 from power plants in seawater to form carbonate ions (CO_3^{2-}) which mix with calcium in the ocean to form a solid cement alternative whilst sequestering CO_2 .

$\text{CO}_2 + \text{H}_2\text{O} \rightarrow \text{H}_2\text{CO}_3 \rightarrow \text{CO}_3^{2-} + 2\text{H}^+$
 $\text{CO}_3^{2-} + \text{Ca}^{2+} \rightarrow \text{CaCO}_3$

PORIFERA

Glass sponge

ANCHOR SPICULES
Around 200 spicules anchor each glass sponge to the sea floor. The below ground morphology forms a strong hold-fast but also allows for considerable pliancy rather than rigidity. This allows for a much more efficient anchoring system in terms of materials. Potential: ANCHORING OFF-SHORE WIND TURBINES

FIBRE OPTICS
The silicon dioxide fibers of the glass sponge transmit light better than fibre optics and are made from benign materials at ambient temperature. Manufacture of fibre-optic cables, in contrast, occurs at high temperatures where the addition of specialized impurities, such as sodium which improves light conduction, is difficult. Glass sponges could help us design flexible (rather than brittle), faster fibre optic cables using far less energy.

HIERARCHICAL STRUCTURE

ARTHROPODA

COLEOPTERA

Darkling Beetle

WATER COLLECTION
Namibian desert beetles collect water from air by using the wind with their tails in the air. Water collects on hydrophilic bumps on the beetle's exoskeleton & then runs through hydrophobic troughs to a collector which runs straight into their mouth.

SELF-FILLING WATER BOTTLES
Self-filling water bottles have been designed using the water collection mechanism used by Namibian desert beetles.

FOG NETS
Namibian desert beetle inspired fog nets could be used to collect drinking & agricultural water in some of the driest parts of the world.

AGRICULTURAL IRRIGATION

CRANIATA

CETACEA

Humpback whale


HYDRODYNAMICS
Humpback whales have long pectoral fins with uneven leading edges. The protuberances of the fins, called tubercles, induce vortices which create extra lift & allow the whale to maintain maneuverability at low speeds.

WIND TURBINES
A new wind turbine blade incorporates tubercles to produce a wind turbine that is 20% more efficient & maintains operation at low speeds.

WIND TURBINE WITH TUBERCLES

BIOMIMICRY OF INVERTEBRATES

BY GEMMA SHEPHERD



CRANIATA

CHONDRICTHYES

Galapagos shark

ANTI-BACTERIAL SKIN
The skin of a shark is covered in tiny scales called dermal denticles which are arranged parallel to water flow. These dermal denticles are covered in microscopic bumps & hollows. The topography makes the surface inhospitable to bacteria & other organisms as it would require too high an energy demand to reside upon.

DERMAL DENTICLES

ANTI-MICROBIAL SURFACES

ANTI-BACTERIAL SURFACE
Inspired by the patterns of shark skin you can produce a polymer topography which prevents bacteria developing on its surface. This reduces the need for anti-bacterial chemical agents & could be very useful in places like hospitals.

ARTHROPODA

HEXAPODA

Termites

TEMPERATURE REGULATION
Termites can keep their mounds at a constant temperature of 29°C-32°C despite the outdoor temperature ranging by 40°C. They do this by constructing sophisticated ventilation systems with porous walls & vents which the termites can open & close.

VENTILATION
A termite inspired Eastgate centre was built in Zimbabwe with a self-regulating ventilation system with openings that promote passive air flow through high thermal capacity building materials. This mechanism could vastly improve buildings' energy efficiency.

ARTHROPODA

CHELICERATA

Spiders

STRONG SILK
Spiders produce different types of liquid silk in their silk glands which are released as a gel through spinnerets on their abdomen. The silk consists of a fibrous core filament covered in other protein adhesives (sericins). The gel is drawn out of the spinnerets as a fine thread stronger than steel & more elastic than latex. The silk also has 'shape memory' & is never soluble in water.

LIGHTWEIGHT MATERIALS
Generating a material which mimics spider silk could have application in the biomedical domain, for sports of military equipment, or any industry looking for a strong, elastic, lightweight & benign material.

MOLLUSK

BIVALVIA

Mussels

ADHESION
Mussels adhere to both wet & dry surfaces by secreting strong, wear-resistant & elastic byssal threads made up of bundles of collagen surrounded by a thin cuticle.

ADHESIVE AMINO ACID
 $\text{= 3,4-dihydroxyphenylalanine}$

CATECHOL FUNCTIONAL GROUP
Forms strong bonds with adjacent molecules & metal ions on solid surface.

NON-TOXIC ADHESIVES
Mussel inspired adhesives can be made by modifying soy proteins so that the amino acids mimic those found in byssal threads. This produces a non-toxic, water resistant adhesive alternative to carcinogenic formaldehyde adhesives.

MOLLUSK

CEPHALOPOD

Cuttlefish

CAMOUFLAGAGE
Cuttlefish have chromatophores (pigment cells) in their skin which can be made visible or invisible using muscles under direct control from neurons from the brain. They also have leucophores & iridophores, pigment cells under the chromatophores which help reflect light. Iridophores use a stack of platelets containing the protein reflectin to produce iridescent reflections whilst leucophores on the bottom reflect almost all incident light. They can also change the texture of their skin using muscle bundles called papillae.

COLOUR & LIGHT
Aside from the use of this mechanism for camouflage, it could inspire the development of energy efficient screens & revolutionize how we colour the world around us.

TISSUE REGENERATION
Platyhelminthes such as Dugesia have the remarkable ability to regenerate full bodies from cut-up segments. Severing body parts sets off a cascade of stem cell signals which activate the creation of the body part that has been lost. Some flatworms even use this as a method of asexual reproduction.

MEDICINE
Researchers analysing the composition of stem cells & proteins in flatworms have discovered a protein critical for the maintenance of pluripotent adult stem cells in both humans & platyhelminthes. Translating our understanding of regeneration in flatworms could guide the development of regenerative medicine in humans.

TARDIGRADA


Water bear / Moss piglet

RESISTANCE
Tardigrades can survive in very extreme conditions. They are resistant to X-rays, can survive in vacuum, high pressures, high temperatures, extreme cold & can survive extreme desiccation. To do this they use a mechanism of cryptobiosis in which they enter an anabolic state, expel 99% of their water & replace it with a sugar which acts as an anti-freeze. The tardigrade then rolls up into a tun & is covered in a layer of wax until normal conditions return.


CRYPTOBIOSIS

PRESERVATION
Understanding cryptobiosis & its preservation mechanisms could open up new possibilities in food preservation & the transport & preservation of organs & other tissue for medical use.


PORIFERA A HEXACTINELLIDA Glass sponge



ANCHOR SPICULES



Around 200 spicules anchor each glass sponge to the sea floor. The below ground morphology forms a strong hold-fast but also allows for considerable pliancy rather than rigidity. This allows for a much more efficient anchoring system in terms of materials. Potential: ANCHORING OFF-SHORE WIND TURBINES




HIERARCHICAL STRUCTURE



The silicon dioxide fibers of the glass sponge transmit light better than fibre optics and are made from benign materials at ambient temperature. Manufacture of fibre-optic cables, in contrast, occurs at high temperatures where the addition of specialized impurities, such as sodium which improves light conduction, is difficult. Glass sponges could help us design flexible (rather than brittle), faster fibre optic cables using far less energy.

CNIDARIAN ANTHOZOA

Stony corals





CORAL GROWTH



Polyps secrete skeletons of calcium carbonate (CaCO_3) in the form of a cup in which the polyp sits. Occasionally the polyps lift up & create a new base above it, elevating the coral upward using carbon dioxide & calcium from the ocean.

CARBON SEQUESTERING CEMENT

Mimicing the mechanism by which corals build thier skeletons could transform th cement industry, which is currently responsible for 8% of total CO_2 emmissions. The cement company Calcera now makes cement by dissolving waste CO_2 from power plants in seawater to form carbonate ions (CO_3^{2-}) which mix with calcium in the ocean to form a solid cement alternative whilst sequestering CO_2 .



ECHINODERMATA HOLOTHUROIDEA Sea cucumber



RIGIDITY & FLEXIBILITY

Sea cucumbers have mutable collagenous tissues which can extensively change their mechanical properties. Their skin can switch between soft, normal & rigid states in the blink of an eye. Their skin contains collagen fibrils which are able to bind strongly or not with other substances around them, switching from an interlinked rigid state to a loose flexible state.

SMART POLYMER

Researchers studying sea cucumber skin have developed a polymer which is flexible in aqueous solvent & becomes rigid when the solvent evaporates. They hope to use this for 'setting' medical devices & responsive brain implants.

SOFT ← NORMAL → HARD
- Hydrogen bonding inhibited - H. bonding activated.



Electrical impulses


CRANIATA CHONDRICHTHYES Galapagos shark




ANTI-BACTERIAL SKIN

The skin of a shark is covered in tiny scales called dermal denticles which are arranged parallel to water flow. These dermal denticles are covered in microscopic bumps & hollows. The topography makes the surface inhospitable to bacteria & other organisms as it would require too high an energy demand to reside upon.

DERMAL
DENTICLES



ANTI-MICROBIAL
SURFACES



ANTI-BACTERIAL SURFACE

Inspired by the patterns of sharkskin you can produce a polymer topography which prevents bacteria developing on its surface. This reduces the need for anti-bacterial chemical agents & could be very useful in places like hospitals.

CRANIATA CETACEA Humpback whale



HYDRODYNAMICS

Humpback whales have long pectoral fins with uneven leading edges. The protuberances of the fins, called tubercles induce vortices which create extra lift & allow the whale to maintain maneuverability at low speeds.

WIND TURBINES

A new wind turbine blade incorporates tubercles to produce a wind turbine that is 20% more efficient & maintains operation at low speeds.



WIND TURBINE WITH TUBERCLES

UROCHORDATA THALIACEA

Salps



PARTICLE CAPTURE

Salps continuously secrete mucus nets which can trap particles smaller than the size of the mesh. They feed by rhythmically pumping water into their oral siphon, through the pharyngeal chamber with the mucus net, & out of their atrial siphon. By pumping water into their bodies they reduce the effects of turbulence so bacteria, viruses & colloidal matter get trapped on the mucus secretion. This mechanism allows salps to survive on a diet of the smallest biological life forms.

WATER FILTRATION


This filtration mechanism could be used to capture & remove sub-micron particles from drinking water.

PLATYHELMINTHES

Flatworms



TISSUE REGENERATION




Platyhelminthes such as *Dugesia* have the remarkable ability to regenerate full bodies from cut up segments. Severing body parts sets off a cascade of stem cell signals which activate the creation of the body part that has been lost. Some flatworms even use this as a method of asexual reproduction.

MEDICINE

Researchers analysing the composition of stem cells & proteins in flatworms have discovered a protein critical for the maintenance of pluripotent adult stem cells in both humans & platyhelminthes. Translating our understanding of regeneration in flatworms could guide the development of regenerative medicine in humans.



MOLLUSK CEPHALOPOD Cuttlefish




CAMOUFLAGE

Cuttlefish have chromatophores (pigment cells) in their skin which can be made visible or invisible using muscles under direct control from neurons from the brain. They also have leucophores & iridophores, pigment cells under the chromatophores which help reflect light. Iridophores use a stack of platelets containing the protein reflectin to produce iridescent reflections whilst leucophores on the bottom reflect almost all incident light. They can also change the texture of their skin using muscle bundles called papillae.

COLOUR & LIGHT

Aside from the use of this mechanism for camouflage, it could inspire the development of energy efficient screens & revolutionize how we colour the world around us.




MOLLUSK BIVALVIA mussels

ADHESION Mussels adhere to both wet & dry surfaces by secreting strong, wear-resistant & elastic byssal threads made up of bundles of collagen surrounded by a thin cuticle.



ADHESIVE AMINO ACID

= 3,4 dihydroxyphenylalanine



CATECHOL FUNCTIONAL GROUP

forms strong bonds with adjacent molecules & metal atoms on solid surface.

NON-TOXIC ADHESIVES

Mussel inspired adhesives can be made by modifying soya proteins so that the amino acids mimic those found in byssal threads.


This produces a non-toxic, water resistant adhesive alternative to carcinogenic formaldehyde adhesives.

TARDIGRADA Water bear / Moss piglet



RESISTANCE

Tardigrades can survive in very extreme conditions. They are resistant to X-rays, can survive in vacuums, high pressures, high temperatures, extreme cold & can survive extreme desiccation. To do this they use a mechanism of cryptobiosis in which they enter an ametabolic state, expel 99% of their water & replace it with a sugar which acts as an anti-freeze. The tardigrade then rolls up into a tun & is covered in a layer of wax until normal conditions return.






PRESERVATION

Understanding cryptobiosis & its preservation mechanism could open up new possibilities in food preservation & the transport & preservation of organs & other tissue for medical use.

ARTHROPODA

COLEOPTERA

Darkling Beetle




WATER COLLECTION

Namibian desert beetles collect water from mist by facing the wind with their tails in the air. Water collects on hydrophilic nanobumps on the beetle's exoskeleton & then runs through hydrophobic troughs to a collector which runs straight into their mouth.

SELF-FILLING

WATER BOTTLES



Self-filling water bottles have been designed using the water collection mechanism used by Namibian desert beetles.

FOG NETS



Namibian desert beetle inspired fog nets could be used to collect drinking & agricultural water in some of the driest parts of the world.



AGRICULTURAL IRRIGATION

ARTHROPODA CHELICERATA Spiders



STRONG SILK

Spiders produce different types of liquid silk in their silk glands which are released as a gel through spinnerets on their abdomen. The silk consists of a fibroin core filament covered in other protein adhesives (sericins). The gel is drawn out of the spinnerets as a fine thread stronger than steel & more elastic than latex. The silk also has 'shape memory' & is never soluble in water.



LIGHTWEIGHT MATERIALS

Generating a material which mimics spider silk could have applications in the biomedical domain, for sports or military equipment, or any industry looking for a strong, elastic, light weight & benign material.

ARTHROPODA

HEXAPODA

Termites



TEMPERATURE REGULATION

Termites can keep their mounds at a constant temperature of 29°C – 32°C despite the outdoor temperature ranging by 40°C . They do this by constructing sophisticated ventilation systems with porous walls & vents which the termites can open & close.

VENTILATION


A termite inspired Eastgate centre was built in Zimbabwe with a self regulating ventilation system with openings that promote passive air flow through high thermal capacity building materials. This mechanism could vastly improve buildings energy efficiency.

ARTHROPODA LEPIDOPTERA Morpho Butterfly



COLOUR EFFECTS

Butterfly wings consist of multi-layered ribbed structures, covered by scales. The colours perceived on the wings are a function of the shape, size & arrangement of the scales. The butterfly wing consists of several strata of pigment & structural scales. which overlap. For example, the iridescent blue of morpho butterflies is produced by structural bottom scales & transparent cover scales.



COLOUR & WINDOWS

By mimicing the colour effects of butterflies we could develop materials of any colour without using any pigment, dyes, water or treatment. Furthermore, this could be used to produce spectrally selective glass which could reduce the need for low emissivity coatings on windows which are usually applied as a seperate material.

Sources - Biomimicry of Invertebrates

Agnarsson, I., Kuntner, M., & Blackledge, T. A. (2010). Bioprospecting finds the toughest biological material: extraordinary silk from a giant riverine orb spider. *PloS one*, 5(9), e11234.

Birenheide, R., Tamori, M., Motokawa, T., Ohtani, M., Iwakoshi, E., Muneoka, Y., ... & Nomoto, K. (1998). Peptides controlling stiffness of connective tissue in sea cucumbers. *The Biological Bulletin*, 194 (3), 253-259.

Bixler, G. D., & Bhushan, B. (2012). Biofouling: lessons from nature. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 370 (1967), 2381-2417.

Boothby, T. C., Tapia, H., Brozena, A. H., Piskiewicz, S., Smith, A. E., Giovannini, I., ... & Goldstein, B. (2017). Tardigrades use intrinsically disordered proteins to survive desiccation. *Molecular cell*, 65(6), 975-984.

Brooks, M. (2008). Do you speak cuttlefish?. *New Scientist*, 198 (2653), 28-31.

Capadona, J. R., Shanmuganathan, K., Tyler, D. J., Rowan, S. J., & Weder, C. (2008). Stimuli-responsive polymer nanocomposites inspired by the sea cucumber dermis. *science*, 319 (5868), 1370-1374.

Chung, K. K., Schumacher, J. F., Sampson, E. M., Burne, R. A., Antonelli, P. J., & Brennan, A. B. (2007). Impact of engineered surface microtopography on biofilm formation of *Staphylococcus aureus*. *Biointerphases*, 2 (2), 89-94.

Constantz, B. R. (1986). Coral skeleton construction: a physiochemically dominated process. *Palaos*, 152-157.

Dizikes, P. (2011, April 21). Out of thick air. Retrieved from <https://news.mit.edu/2011/fog-harvesting-0421>.

Fish, F. E., & Battle, J. M. (1995). Hydrodynamic design of the humpback whale flipper. *Journal of Morphology*, 225 (1), 51-60.

Getting a Head: How Worms Regenerate Lost Tissue. (2019). Retrieved from <https://www.livescience.com/38405-how-flatworms-regrow-their-heads.html>.

Hamilton III, W. J., Henschel, J. R., & Seely, M. K. (2003). Fog collection by Namib Desert beetles: correspondence. *South African Journal of Science*, 99 (3-4), 181.

Hori, I., & Kishida, Y. (1998). A fine structural study of regeneration after fission in the planarian *Dugesia japonica*. *Hydrobiologia*, 383 (1-3), 131-136.

Lee, H., Dellatore, S. M., Miller, W. M., & Messersmith, P. B. (2007). Mussel-inspired surface chemistry for multifunctional coatings. *science*, 318 (5849), 426-430.

Monn, M. A., Weaver, J. C., Zhang, T., Aizenberg, J., & Kesari, H. (2015). New functional insights into the internal architecture of the laminated anchor spicules of *Euplectella aspergillum*. *Proceedings of the National Academy of Sciences*, 112 (16), 4976-4981.

Ocko, S. A., King, H., Andreen, D., Bardunias, P., Turner, J. S., Soar, R., & Mahadevan, L. (2017). Solar-powered ventilation of African termite mounds. *Journal of Experimental Biology*, 220(18), 3260-3269.

Pawlyn, M. (2019). *Biomimicry in architecture*. Routledge.

Plantkart. (2019). Calera MAP cement-making process. Retrieved from <https://asknature.org/idea/calera-map-cement-making-process/#.Xd1vYJpKiM8>.

Ralevski, A., Larson, S., Sherry, Pechlivanoglou, G., & Greiser, T. (2019). Tubercle Technology blades. Retrieved from <https://asknature.org/idea/tubercle-technology-blades/>.

Romero, A. (2016, September 6). Structural Color: When light transforms into art. Retrieved from <https://annabelromero.com/opuntiavisual/2016/9/4/structural-color-when-light-transforms-into-art>.

Sullivan, T., & Regan, F. (2011). The characterization, replication and testing of dermal denticles of *Scyliorhinus canicula* for physical mechanisms of biofouling prevention. *Bioinspiration & biomimetics*, 6 (4), 046001.

Sundar, V. C., Yablon, A. D., Grazul, J. L., Ilan, M., & Aizenberg, J. (2003). Fibre-optical features of a glass sponge. *Nature*, 424 (6951), 899.

Sutherland, K. R., Madin, L. P., & Stocker, R. (2010). Filtration of submicrometer particles by pelagic tunicates. *Proceedings of the National Academy of Sciences*, 107 (34), 15129-15134.

Gemma Shepherd

Team, A. N. (2019). Mechanically adaptive polymer nanocomposites. Retrieved from <https://asknature.org/idea/mechanically-adaptive-polymer-nanocomposites/#.XeJ-4JpKiM8>.

Team, A. N. (2019.). Eastgate Centre. Retrieved from <https://asknature.org/idea/eastgate-centre/>.

Ternaux, E., Ternaux, M., & Ternaux, J. P. (2012). *Industry of Nature: another approach to ecology*. Frame Publishers.

Vukusic, P. (2006). Structural colour in Lepidoptera. *Current Biology*, 16 (16), R621-R623.

Weaver, J. C., Aizenberg, J., Fantner, G. E., Kisailus, D., Woesz, A., Allen, P., ... & Fratzl, P. (2007). Hierarchical assembly of the siliceous skeletal lattice of the hexactinellid sponge *Euplectella aspergillum*. *Journal of structural biology*, 158 (1), 93-106.

Woolley-Barker, T., Harun, Y., Brooks, M., Ralevski, A., Sherry, Greiser, T., & Benyus, J. (2002, January 17). Rapid color change used for protection : Cuttlefish. Retrieved from <https://asknature.org/strategy/rapid-color-change-used-for-protection/>.